

Nipah Virus Panic in Kerala

What is the issue?

 $n\n$

\n

- The lethal Nipah Virus has been spreading in northern Kerala, creating a health emergency and widespread panic.
- The Zoonotic virus is suspected to have spread to humans from infected bats, and actions to curtail the spread of infections are currently underway.

 $n\n$

How is the current episode panning out?

 $n\$

\n

- \bullet Nipah spreads from fruit bats to humans and other animals, mainly through bad dropping or bodily remains and then spreads laterally within a species. \n
- But less than 1% of the fruit bats are estimated be infected with Nipah virus and it is even rarer for it to infect humans.
- It is indeed highly contagious within humans, and the current episode in Kerala even killed a nurse who was treating another Nipah patient.
- The outbreak has already cost the lives of 10 people out of 12 confirmed cases in Kozhikode and Malappuram of Kerala.
- Notably, the previous serious outbreak in India was in 2001, when the Siliguri district of West Bengal recorded 42 Nipah related deaths.
- The disease has a high mortality rate, but experts stress that there is no reason to panic because Nipah outbreaks have always been generally localised.

 $n\n$

What are the symptoms of Nipah?

 $n\n$

۱n

- \bullet The classical symptom is acute and rapidly progressive encephalitis (brain inflammation and pain) with or without respiratory involvement. \n
- Other more noticeable symptoms include non-productive cough during the early part of the disease.
- Nipah encephalitis comes with 3-14 days of fever and headache, followed by drowsiness, disorientation and mental confusion.
- \bullet Acute encephalitis progresses to coma within 24-48 hours. $\ensuremath{\backslash n}$

 $n\n$

How can the spread of Nipah be contained?

 $n\n$

۱n

 \bullet Virologists working on the ground have asserted that early diagnosis has helped in containing the spread in the current case. \n

 $n\n$

\n

- Timely laboratory confirmation and aggressive tracing of those who came in contact with patient can contain spread of the virus.
- This is very important because there is no effective specific treatment for the infection and hospitalisation is only to support our bodily immune systems.
- Standard infection control practices (like washing and sanitising) and proper barrier nursing techniques are also critical for curtailing spread.

 $n\n$

 $n\n$

Source: Indian Express

\n

