



Understanding the TB Challenge

What is the issue?

\n\n

\n

- India has a high prevalence of TB with an annual incidence of 28 lakhs or 27% of the total global TB incidences.

\n

- Hence, India is currently the TB capital of India and to outsmart the disease, India must intercept - infection, progression and transmission.

\n

\n\n

What is TB?

\n\n

\n

- Tuberculosis (TB) is a disease caused when one gets infection with TB bacilli.

\n

- It has the unique character of mimicking other diseases and hence confuses doctors, which delays diagnosis and further treatment.

\n

- Notably, the common symptom associated with Cough and blood in sputum occurs only in **lung TB** and there are others like **Brain TB, Pelvic TB** etc...

\n

- TB can affect the lungs, brain, bones, joints, the liver, intestines or for that matter any organ and can progress slowly or kill in weeks.

\n

- There are 3 distinct stages in TB - infection, progression, transmission.

\n

- **Infection** - Infection occurs when TB bacilli are inhaled and the Bacilli may stay in the lungs or travel to other organs.

\n

- Once infected, the bacillus persists lifelong inside an organism's body, but lies dormant - This phase is "latent TB".

- \n
- This can be diagnosed by a '**Tuberculin Skin Test**' (TST) and cumulatively, 40% to 70% of us are estimated to be living with latent TB.
- \n
- **Progression** - From this dormant bacterium pool, a few germs slowly lead to the progression of disease in a time span of anywhere between 5 - 30 years.
- \n
- Hence, the TB disease sets in only when bacilli become active and starts multiplying, and this phase is called "active TB".
- \n
- **Transmission** - Only when active TB affects the lungs do bacilli find an exit route to the atmosphere, which is the only mode of transmission.
- \n

\n\n

What is the current status of TB in India?

\n\n

- \n
- The incidence rate of tuberculosis (TB) in India is estimated at 200-300 cases per lakh population per year, whereas it is only 5 per lakh in Europe.
- \n
- Revised National TB Control Programme (RNTCP) provides for free diagnosis and treatment, but it doesn't track incidences and there are no targets.
- \n
- Estimates are that every day 1,200 Indians die of TB, a calamity rate unrivalled by any other disease and a clear indication that control efforts are failing.
- \n
- Hence, this calls for a more robust and specific health interventions to foster a deliberate reduction of incidence to as low as practicable.
- \n

\n\n

What are the challenges?

\n\n

- \n
- While curative treatment is the patient's urgent need, it will not control the spread of TB as germs already gets spewed due to delayed diagnosis.
- \n
- As cough is a very common symptom, TB isn't suspected until other treatments have failed, which delays targeted treatment.
- \n

- Inclination to self-medication through pharmacies, where patients don't see a doctor until things get complicated is another worry in India.
\n
- Additionally, according to RNTCP guidelines, testing is done only after two weeks of consistent cough, which results in the loss of precious lead time.
\n
- Also, 70% of people seek treatment in the private sector and as the mandatory reporting system for TB isn't rigorous - incidences might be underestimated.
\n
- Hence, strengthening public-private sector health partnership and establishing efficient primary health care services is essential for early diagnosis of TB.
\n

\n\n

What is the way forward?

\n\n

- Dealing with TB requires mass initiatives - health professionals, policy planners and administrators and the public must come together to solve it.
\n
- **Prevention** - Public education on TB and its prevention must replace ignorance and misconceptions that are presently prevalent.
\n
- While dealing with the disease after onset is difficult, it is easier to prevent transmission and infection by healthy public etiquettes.
\n
- TB cases can be greatly reduced if basic health sensitiveness of not spitting in public and "mouth covered cough and sneeze practices" are adopted.
\n
- Notably, a person with "**Lung TB**" disseminates TB bacilli over several weeks and by the time dissemination stops, it would've already affected many others.
\n
- To block transmission, treatment should begin as soon as symptoms show up, which will effectively bring down incidences.
\n
- **Treating latent TB** - Drug treatment of "latent TB" is an option and it has been recommended for all children in the 5-10 age group to be screened for **TST**.
\n
- Treatment of latent TB will prevent its progression to active TB and

consequently bring down the “Annual Risk of Tuberculous Infection” (ARTI).

\n

- While these will take as much as 2 to 3 decades, this is the only way to achieve sustained reduction of TB incidences and a start has to be made now.

\n

- Any further delay can be catastrophic as the TB microbes are increasingly becoming multi drug resistant.

\n

\n\n

\n\n

Source: The Hindu

\n\n

\n



IAS PARLIAMENT

Information is Empowering

A Shankar IAS Academy Initiative